

Lettuce corals overgrowing tube sponges at St. Eustatius, Dutch Caribbean

Jaaziel E. García-Hernández^{1,2} · Godfried W. N. M. van Moorsel³ · Bert W. Hoeksema^{2,4}

Received: 23 December 2015 / Revised: 8 February 2016 / Accepted: 16 February 2016 / Published online: 11 March 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Sponges and corals are well-known competitors for space on coral reefs. Only a few documented observations are known of sponge-coral interactions in which the latter dominate the former. In the Caribbean, reef-dwelling sponges are generally abundant and have a reputation of being aggressive competitors for space. The present study is exceptional, because it reports on corals overgrowing sponges.

Keywords Competition for space · Coral reefs · Interspecific dominance · Sponge-coral interactions

Within Caribbean coral reef ecosystems, interactions between corals and sponges are well documented, usually with sponges outcompeting corals in direct competition for growth space (Aerts and Van Soest 1997). Some of these sponges are able to produce toxic compounds that

may harm and kill scleractinian corals, as demonstrated in interactions between the liver sponge *Plakortis halichondrioides* (Wilson, 1902) and the lettuce coral *Agaricia lamarcki* (Milne Edwards and Haime, 1851) (Porter and Targett 1988). Other Caribbean sponges may harm corals by overgrowing them (Loh et al. 2015) or by excavating their skeleton (López-Victoria et al. 2006). Documented examples of sponges becoming partly or completely submissive to corals are exceptional. These are known, for instance, from the Indo-Pacific, involving fast-growing encrusting parts of foliaceous *Montipora* corals (Elliott et al. 2015) and free-living corals that have dropped and settled on phototrophic sponges (Hoeksema et al. 2014).

During a marine biodiversity survey at St. Eustatius (Lesser Antilles) in June 2015, a colony of *Agaricia agaricites* (Linnaeus, 1758) was found at 19 m depth overgrowing a brown tube sponge, *Agelas conifera* (Schmidt, 1870) (Fig. 1a). The sponge showed signs of tissue damage and settlement of red filamentous algae (Fig. 1b). This observation was unexpected, since *A. conifera* is regarded as a chemically defended sponge which is able to deter fish predation, inhibit growth of other sponges, and prevent fouling organisms from colonizing it (Assmann et al. 2004). This interaction was observed only once during the whole expedition (40 survey sites), whereas both species were found to be co-occurring at 21 sites. In addition, a sunray lettuce coral, *Helioseris cucullata* (Ellis and Solander, 1786), of the same family (Agariciidae) was found to overgrow

Communicated by R. Vonk

✉ Bert W. Hoeksema
bert.hoeksema@naturalis.nl

¹ Department of Biology, University of Puerto Rico at Mayagüez, P.O. Box 9000, Mayagüez, PR 00681, USA

² Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

³ Ecosub, P.O. Box 126, 3940 AC Doorn, The Netherlands

⁴ Institute of Biology Leiden, Leiden University, PO Box 9505, 2300 RA Leiden, The Netherlands




Fig. 1 Coral–sponge interactions at St. Eustatius: **a** *Agaricia agaricites* coral overgrowing part of a tube sponge, *Agelas conifera*; **b** same sponge showing tissue damage and algal settlement (arrows); **c** *Helioseris cucullata* coral partly overgrowing an *A. conifera* sponge. Scale bars: 1 cm

an *A. conifera* tube sponge, with both species reported as co-occurring at 11 sites (Fig. 1c).

The ability of agariciid corals to overgrow chemically defended sponges is worth pursuing. Additional field studies in other Caribbean reefs are needed to confirm whether such interactions are more common than presently assumed.

Acknowledgments Fieldwork participation of the first author was funded through a Martin Fellowship from Naturalis Biodiversity Center. Logistic support was provided by St. Eustatius Marine Parks (STENAPA), Caribbean Netherlands Science Institute (CNSI) and Scubaqua Dive Center. We are grateful to three anonymous reviewers for their constructive comments.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Aerts LAM, van Soest RWM (1997) Quantification of sponge/coral interactions in a physically stressed reef community, NE Colombia. *Mar Ecol Prog Ser* 148:125–134
- Assmann M, Lichte E, Köck M (2004) Multiple defensive roles for Bromopyrrole alkaloids from Caribbean *Agelas* sponges. *Boll Mus Ist Biol Univ Genova* 68:187–193
- Elliott J, Patterson M, Vitry E, Summers N, Miteron C (2015) Morphological plasticity allows coral to actively overgrow the aggressive sponge *Terpios hoshinota* (Mauritius, Southwestern Indian Ocean). *Mar Biodiv*. doi:10.1007/s12526-015-0370-4
- Hoeksema BW, Dekker F, de Voogd NJ (2014) Free-living mushroom corals strike back by overtopping a coral-killing sponge. *Mar Biodiv* 44:3–4
- Loh TL, McMurray SE, Henkel TP, Vicente J, Pawlik JR (2015) Indirect effects of overfishing on Caribbean reefs: sponges overgrow reef-building corals. *Peer J* 3, e901
- López-Victoria M, Zea S, Weil E (2006) Competition for space between encrusting excavating Caribbean sponges and other coral reef organisms. *Mar Ecol Prog Ser* 312:113–121
- Porter JW, Targett NM (1988) Allelochemical Interactions between sponges and corals. *Biol Bull* 175:230–239